

THIS RECOMMENDATION IS MADE AS PERMITTED UNDER FIFRAL SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.



2(ee) Recommendation

FOR DISTRIBUTION AND USE ONLY IN OREGON

THIS PRODUCT BULLETIN CONTAINS NEW OR ADDITIONAL DIRECTIONS FOR USE WHICH ARE RECOMMENDED BY FMC CORPORATION UNDER FIFRA SECTION 2(ee)

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODCUT IN A MANNER INCONSISTENT WITH ITS LABELING

CROP	PEST	RATE OF APPLICATION
Snap Beans	White Mold (<i>Schlerotinia sclerotiorum</i>) Gray Mold (<i>Botrytis cinerea</i>)	Rovral Fungicide: 24 fl. oz/A + Topsin 4.5 Flowable: 22 fl. oz/A

DIRECTIONS FOR USE:

Apply using ground equipment with a spray pressure of 50-100 PSI using a three nozzle per row boom arranged with line directly over the row and a drop on each side of the row. Application may also be made by air or chemigation.

TIMING OF APPLICATION:

For high pressure environments begin applications at 10% bloom and follow with sequential treatments 5-7 days later up to peak bloom.

USE RESTRICTIONS:

There is a maximum of two Rovral applications per season, with the last application made no later than peak bloom. Do not apply within 14 days of harvest.

Do not allow foraging for 14 days after the last application.

Do not feed snap bean hay to livestock.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS ON THE EPA REGISTERED ROVRAL® 4 FLOWABLE FUNGICIDE LABEL (EPA REG. NO. 279-9564) MUST BE FOLLOWED. THIS RECOMMENDATION SHOULD BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.



FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia, PA 19103

Rovral and FMC - FMC Trademarks
© 2014 FMC Corporation
Topsin 4.5 Flowable - Trademark of
Nippon Soda Company Ltd.